

# Comment on “Three Extra Mirror or Sequential Families: Case for a Heavy Higgs Boson and Inert Doublet”

M. Sahin<sup>1</sup>, S. Sultansoy<sup>1,2</sup>, G. Unel<sup>3</sup>

<sup>1</sup>TOBB University of Economics and Technology, Ankara, Turkey

<sup>2</sup>ANAS Institute of Physics, Baku, Azerbaijan

<sup>3</sup>University of California at Irvine, Irvine, CA, USA

In a recent Letter [1], Martinez *et al.* have argued that the existence of extra three families is still perfectly in accord with the standard model (SM), as long as an additional Higgs doublet is also present. Here we claim that the mentioned accordance can be obtained with less reliance on BSM physics like additional Higgs doublets, but benefiting from Majorana neutrinos is sufficient. For the remainder of the discussion we will denote SM with  $n$  families as SM $n$  ( $n = 3, 4, 5, 6$ ).

Let us remember that while first papers on the SM did not contain right-handed components of the neutrinos, their inclusion is necessary according quark-lepton symmetry ( $\nu_R$  is counterpart of  $u_R$ ). Moreover, if there are no any arguments (like exact conservation of lepton charge) forbidding Majorana mass terms for  $\nu_R$ , these terms should be taken into account. Of course, this leads to some complication (such as the  $2n \times 2n$  mass matrix in neutrino sector instead of an  $n \times n$  one) but, simultaneously, provides small masses for neutrinos from the first three families via see-saw mechanism. Therefore, Majorana nature for the SM neutrinos seems more natural than Dirac one.

The widely accepted method for measuring the compatibility of a model with additional fermions and the electroweak (EW) precision data is utilization of the oblique parameters [2]. The rather tedious calculation of  $S$  and  $T$  parameters is usually computerized. Among similar computer programs, OPUCEM [3] is the only tool which can include Majorana neutrinos in the calculations. In the simple SM4 case, the inclusion of Majorana neutrinos essentially enlarges the parameter space, resulting in a large number of SM4 mass points which are in better accordance with the data compared to SM3 [4].

Table 1: Two Majorana type SM6 mass points (mass in GeV). First point (P1) with  $m_h=115\text{GeV}$  and  $|\sin\theta_{34}|=0.08$ , leading to  $S=0.082$  and  $T=0.121$  is on the left and the second point (P2) with  $m_h=600\text{GeV}$  and  $|\sin\theta_{34}|=0.06$ , leading to  $S=0.194$  and  $T=0.203$  is on the right .

	4th f.	5th f.	6th f.	4th f.	5th f.	6th f.
$m_U$	430	565	550	430	565	550
$m_D$	385	450	500	385	450	500
$m_\nu$	45	45	450	45	45	450
$m_E$	570	580	600	590	580	600
$m_N$	2600	2800	3200	2600	2800	3200

Concerning SM5 and SM6 OPUCEM analysis show that in the Majorana case there are a lot of mass points




Figure 1: Majorana type SM6 points with low (115 GeV, dark star) and high (600 GeV, light star) Higgs masses as compared to EW precision data error ellipses and the SM3 prediction with  $m_h = 115$  GeV and  $m_t = 173$  GeV (cross).




Figure 2: Dirac type SM6 points with low (115 GeV, dark star) and high (600 GeV, light star) Higgs masses as compared to EW precision data error ellipses and the SM3 prediction with  $m_h = 115$  GeV and  $m_t = 173$  GeV (cross).

in  $1\sigma$  agreement with the precision EW data. For illustration we present two SM6 points in S-T plane in Fig. 1 with corresponding parameters presented in Table 1 (where  $\nu$  and  $N$  denote light and heavy Majorana neutrinos, respectively). It is seen that SM6 point with  $m_h = 115$  GeV is closer to central value than SM3 point.

Keeping in mind that sometimes Majorana neutrinos are interpreted as BSM physics, OPUCEM results for Dirac neutrinos are presented in Fig. 2 (corresponding parameters are presented in Table 2). It is seen that even with Dirac neutrinos precision electroweak data allows three additional SM families within  $2\sigma$ ! Therefore, results of [1] are neither fascinating nor surprising.

Finally, let us mention that there are phenomenological arguments against fifth (and certainly sixth) SM family (see, e.g. [4] and references therein). Nevertheless, the fact that even SM6 is not contradict to present EW data

Table 2: Two Dirac type SM6 mass points (mass in GeV). First point with  $m_h = 115$  GeV and  $|\sin\theta_{34}| = 0.00$ , leading to  $S = 0.25$  and  $T = 0.26$  is on the left and second point with  $m_h = 600$  GeV and  $|\sin\theta_{34}| = 0.01$ , leading to  $S = 0.31$  and  $T = 0.31$  is on the right.

	4th f.	5th f.	6th f.	4th f.	5th f.	6th f.
$m_U$	530	550	580	580	580	590
$m_D$	520	550	580	580	580	590
$m_\nu$	50	50	50	50	50	50
$m_E$	110	130	120	110	140	180

is quite intriguing.

## References

- [1] H. Martinez, A. Melfo, F. Nesti, and G. Senjanovic, Phys. Rev. Lett. **106**, 191802 (2011).
- [2] M. E. Peskin and T. Takeuchi, Phys. Rev. D **46**, 381 (1992).
- [3] O. Cobanoglu, E. Ozcan, S. Sultansoy, and G. Unel, Computer Physics Communications **182**, 1732 (2011); OPUCEM project at CEDAR Hepforge repository: <http://projects.hepforge.org/opucem/>.
- [4] M. Sahin, S. Sultansoy, and S. Turkoz, Phys. Rev. D **83**, 054022 (2011).